arXiv:0807.4167v3 [hep-ph] 12 Sep 2008

MAN/HEP/2008/20
arXiv:0807.4167
July 2008

Supersymmetric Higgs Singlet Effects on B-Meson FCNC Observables at Large $\tan\beta$

Robert N. Hodgkinson[a,b] and Apostolos Pilaftsis[a]

[a] *School of Physics and Astronomy, University of Manchester*
Manchester M13 9PL, United Kingdom

[b] *High Energy Physics Group, Dept. ECM, Univ. de Barcelona*
Av. Diagonal 647, E-08028 Barcelona, Catalonia, Spain

ABSTRACT

Higgs singlet superfields are usually present in most extensions of the Minimal Supersymmetric Standard Model (MSSM) that address the μ-problem, such as the Next-to-Minimal Supersymmetric Standard Model (NMSSM) and the Minimal Nonminimal Supersymmetric Standard Model (MNSSM). Employing a gauge- and flavour- covariant effective Lagrangian formalism, we show how the singlet Higgs bosons of such theories can have significant contributions to B-meson flavour-changing neutral current (FCNC) observables for large values of $\tan\beta \gtrsim 50$ at the 1-loop level. Illustrative results are presented including effects on the B_s and B_d mass differences and on the rare decay $B_s \to \mu^+\mu^-$. In particular, we find that depending on the actual value of the lightest singlet pseudoscalar mass in the NMSSM, the branching ratio for $B_s \to \mu^+\mu^-$ can be enhanced or even suppressed with respect to the Standard Model prediction by more than one order of magnitude.

1 Introduction

Supersymmetry (SUSY), softly broken at the electroweak (EW) or TeV scale, provides one of the most self-consistent frameworks that enables one to technically address the so-called gauge hierarchy problem. In particular, SUSY protects the EW scale from quantum corrections that could be induced by possible new dynamics at the Grand Unified Theory (GUT) or Planck scale. Whilst the exact mechanism of SUSY breaking remains unknown so far, supersymmetric theories do in general contain many sources of FCNC's amongst the soft SUSY-breaking terms. In order to maintain agreement with experimental data, the hypothesis of Minimal Flavour Violation (MFV) is frequently assumed, where the well-known Glashow–Iliopoulos–Maiani (GIM) mechanism can be naturally implemented [1]. Within the MFV framework, all phenomena of flavour and CP violation in the theory originate from the Cabbibo-Kobayashi-Maskawa (CKM) mixing matrix $\mathbf{V}$ [2], such that all FCNC interactions vanish in the limit $\mathbf{V} \to \mathbf{1}_3$.

The FCNC effects on B- and K-meson observables mediated by Higgs bosons have been well studied in the MSSM [3–7]. At large values of $\tan\beta = v_2/v_1$, the ratio of the Higgs doublet vacuum expectation values (VEVs), the couplings of the Higgs bosons to down-type quarks becomes enhanced and 1-loop threshold effects can make a significant contribution to FCNC processes, such as the $B_s - \bar{B}_s$ and $B_d - \bar{B}_d$ mixings, and the leptonic decays $B_{s,d} \to \mu^+\mu^-$.

On the other hand, along with the supersymmetric trilinear Yukawa interactions, the superpotential of the MSSM contains a bilinear SUSY term, $\mu\hat{H}_1\hat{H}_2$, that couples the two Higgs-doublet superfields. This μ-term is phenomenologically necessary in order to break the chiral Peccei-Quinn (PQ) symmetry $U(1)_{\rm PQ}$ and to provide a mass for the as-yet experimentally unobserved higgsinos. The μ-parameter has dimensions of mass and is singlet under the Standard Model (SM) gauge group. It can therefore be driven by supergravity quantum effects to the mass scales $M_{\rm GUT}$ or $M_{\rm Planck}$. However, successful electroweak symmetry breaking in the MSSM implies that μ must be small, close to $M_{\rm SUSY}$, namely at the scale at which SUSY is softly broken. This theoretical difficulty is often referred to as the μ-problem of the MSSM [8, 9].

The μ-problem may be resolved by allowing an effective μ parameter to be generated at the SUSY breaking scale, as the VEV of (the scalar component of) an additional gauge singlet Higgs superfield, $\hat{S}$. The Higgs bilinear term of the MSSM superpotential is replaced with the trilinear coupling $\lambda\hat{S}\hat{H}_1\hat{H}_2$. In its simplest form, such a model re-introduces the undesirable $U(1)_{\rm PQ}$ symmetry of the superpotential, leading to experimentally excluded electroweak-scale axions. Several mechanisms for breaking the $U(1)_{\rm PQ}$ symmetry have been discussed in the literature [9] and lead to phenomenologically distinct models, such as the Next-to-Minimal Supersymmetric Standard Model (NMSSM) [10] and the Minimal

Non-minimal Supersymmetric Standard Model (MNSSM) [11].

An interesting limit for the NMSSM and the MNSSM is the so-called PQ symmetric limit. If $U(1)_{\rm PQ}$ is weakly broken, then the lightest CP-odd Higgs field A_1 is predominantly singlet and very light, essentially becoming the pseudo-Goldstone boson of this symmetry. In this case, it is therefore natural to expect that a pseudoscalar particle A_1 with a mass of a few GeV may have eluded detection so far. Production of such light, highly singlet pseudoscalars within the context of the NMSSM has been considered through the decays of SM-like Higgs fields [12], in associated production with charginos [13] and in rare decays of Upsilon mesons [14].

In the NMSSM and the MNSSM, the singlet Higgs fields have no tree-level couplings to the Standard Model (SM) fermions or gauge bosons. It has long been known that 1-loop threshold corrections can produce significant non-holomorphic Yukawa couplings in the MSSM at large $\tan\beta$ [15–17]. Most recently, it has been realized [18] that analogous threshold corrections produce sizeable radiative Yukawa couplings even for the singlet Higgs bosons in minimal extensions of the MSSM.

In this paper we consider the effects of FCNC Yukawa couplings of the singlet Higgs bosons within the MFV framework. Such effects have been considered in [19, 20] within the NMSSM in the limit of a light pseudoscalar singlet. The scope of our paper goes well beyond these studies, both analytically, as our effective Lagrangian approach includes a resummation of $\tan\beta$ enhanced terms, but also phenomenologically, as we discuss the implications of light singlets at or below the electroweak scale in both the NMSSM and the MNSSM.

The structure of our paper is as follows: in Section 2 we present a manifestly gauge- and flavour-covariant effective Lagrangian framework for the calculation of FCNC vertices between the Higgs bosons and fermions. Section 3 summarises the relevant analytic results regarding FCNC B-meson observables. Numerical results for various FCNC processes are presented in Section 4, for relevant scenarios in both the MNSSM and the NMSSM. Our conclusions are given in Section 5.

2 Effective Lagrangian Formalism

In this section we derive the form of the manifestly gauge- and flavour-covariant effective Lagrangian for Higgs boson interactions with fermions. Our conventions and notations here closely follow those of [7]. The effective Lagrangian describing the down-type quark self-energy transition $Q^0_{jL} \to d^0_{iR}$ may be written in a gauge-symmetric and flavour-covariant form as

$$-\mathcal{L}^d_{\rm eff}[\Phi_1,\Phi_2,S] = \bar{d^0_{iR}}\left(\mathbf{h}_d\Phi_1^\dagger + \Delta\mathbf{h}_d[\Phi_1,\Phi_2,S]\right)_{ij} Q^0_{jL} \;+\; \text{H.c.}\,, \tag{2.1}$$

where $\Phi_{1(2)}$ are the scalar components of the Higgs doublet superfields giving masses to the down-type (up-type) quarks respectively[1], S is the scalar component of the Higgs singlet superfield and $\mathbf{h}_d$ is the 3×3 down-type Yukawa coupling matrix. In (2.1) the first term is the tree level contribution, whilst $\Delta\mathbf{h}_d$ is a 3×3 matrix which is a Coleman–Weinberg effective functional [21] of the background Higgs fields $\Phi_{1,2}$ and S. Observe that the 1-loop effective functional $\Delta\mathbf{h}_d$ has the same flavour- and gauge-transformation properties as $\mathbf{h}_d\,\Phi_1^\dagger$. Typical Feynman graphs that contribute to $\Delta\mathbf{h}_d$ to leading order in the Higgs fields Φ_2 and S are displayed in Fig. 1. In detail, the analytic form of $\Delta\mathbf{h}_d$ is

$$\begin{aligned}(\Delta\mathbf{h}_d)_{ij} \;=\; & \int\frac{d^nk}{(2\pi)^ni}\left[P_L\frac{2C_F\,g_3^2M_3^*}{k^2-|M_3^2|}\left(\frac{1}{k^2\mathbf{1}_{12}-\widetilde{\mathbf{M}}^2}\right)_{\tilde D_i\tilde Q_j^\dagger}\right.\\
&+P_L\left(\frac{1}{\not k\mathbf{1}_9-\mathbf{M}_CP_L-\mathbf{M}_C^\dagger P_R}\right)_{\tilde H_d\tilde H_u}P_L\,(\mathbf{h}_d)_{il}\left(\frac{1}{k^2\mathbf{1}_{12}-\widetilde{\mathbf{M}}^2}\right)_{\tilde Q_l\tilde U_k^\dagger}(\mathbf{h}_u)_{kj}\\
&+P_L\left(\frac{1}{\not k\mathbf{1}_9-\mathbf{M}_CP_L-\mathbf{M}_C^\dagger P_R}\right)_{\tilde H_d\tilde B}P_L\,(\mathbf{h}_d)_{il}\left(\frac{1}{k^2\mathbf{1}_{12}-\widetilde{\mathbf{M}}^2}\right)_{\tilde Q_l\tilde Q_j^\dagger}(\sqrt{2}g_1)\\
&\left.+\sum_{k=1}^3P_L\left(\frac{1}{\not k\mathbf{1}_9-\mathbf{M}_CP_L-\mathbf{M}_C^\dagger P_R}\right)_{\tilde H_d\tilde W^k}P_L\,(\mathbf{h}_d)_{il}\left(\frac{1}{k^2\mathbf{1}_{12}-\widetilde{\mathbf{M}}^2}\right)_{\tilde Q_l\tilde Q_j^\dagger}\left(\frac{g_2\tau^k}{\sqrt{2}}\right)\right]\,.\end{aligned}\tag{2.2}$$

Here $n=4-2\epsilon$ is the number of dimensions in dimensional regularisation (DR), $\mathbf{1}_N$ is the $N\times N$-dimensional identity matrix, $\tau^{1,2,3}$ are the Pauli matrices, $P_{L(R)}=\frac{1}{2}[1-(+)\gamma_5]$ are the standard chirality projection operators and C_F is the quadratic Casimir invariant of QCD in the fundamental representation, i.e. $C_F=4/3$. Moreover, we denote the $U(1)_Y$, $SU(2)_L$ and $SU(3)_c$ gauge couplings by $g_{1,2,3}$ and the soft SUSY-breaking gluino mass by M_3. Finally, $\widetilde{\mathbf{M}}^2$ and $\mathbf{M}_C$ are 12×12- and 9×9-dimensional matrices which describe the squark and chargino-neutralino mass matrices, respectively, in the background of non-vanishing Higgs fields $\Phi_{1,2}$ and S.

We express the 9×9 chargino-neutralino mass matrix $\mathbf{M}_C$ in the Weyl basis

$$\left(\widetilde B,\,\widetilde W^{1,2,3},\,\widetilde H_u,\,\widetilde H_d,\,\tilde s\right),$$

where $\widetilde H_{u,d}$ are $SU(2)_L$ doublets $\widetilde H_u=\left(\tilde h_u^+,\tilde h_u^0\right)^T$, $\widetilde H_d=\left(\tilde h_d^0,\tilde h_d^-\right)^T$ and $\tilde s$ is the singlino Higgs field. In the NMSSM, $\mathbf{M}_C$ is given by

$$\mathbf{M}_C[\Phi_1,\Phi_2,S]=\begin{pmatrix}M_1 & 0 & -\frac{1}{\sqrt2}g_1\Phi_2^\dagger & \frac{1}{\sqrt2}g_1\Phi_1^T(i\tau_2) & 0\\ 0 & M_2\mathbf{1}_3 & \frac{1}{\sqrt2}g_2\Phi_2^\dagger\tau_i & -\frac{1}{\sqrt2}g_2\Phi_1^T(i\tau_2)\tau_i & 0\\ -\frac{1}{\sqrt2}g_1\Phi_2^* & \frac{1}{\sqrt2}g_2\tau_i^T\Phi_2^* & \mathbf{0}_2 & \lambda S(i\tau_2) & -\lambda\Phi_1^*\\ -\frac{1}{\sqrt2}(i\tau_2)g_1\Phi_1 & \frac{1}{\sqrt2}g_2\tau_i^T(i\tau_2)\Phi_1 & -\lambda S(i\tau_2) & \mathbf{0}_2 & -\lambda(i\tau_2)\Phi_2\\ 0 & 0 & -\lambda\Phi_1^\dagger & \lambda\Phi_2^T(i\tau_2) & 2\kappa S\end{pmatrix},\tag{2.3}$$

[1]Here we adopt the convention for the Higgs doublets: $H_u\equiv\Phi_2$, $H_d\equiv i\tau_2\Phi_1^*$, where τ_2 is the usual Pauli matrix.




Figure 1: *Gauge- and flavour-invariant self-energy graphs for the down-type quarks to leading order in the Higgs fields* Φ_2 *and* S*.*

where $M_{1,2}$ are the bino and wino soft SUSY-breaking masses and κ is the trilinear singlet coupling of the NMSSM superpotential; the corresponding matrix for the MNSSM is obtained by setting $\kappa = 0$. The 12×12 squark mass matrix $\widetilde{\mathbf{M}}^2$ is identical in both models. More explicitly, it is given by

$$\widetilde{\mathbf{M}}^2[\Phi_1, \Phi_2, S] = \begin{pmatrix} \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{Q}^\dagger \tilde{Q}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{Q}^\dagger \tilde{U}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{Q}^\dagger \tilde{D}} \\ \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}^\dagger \tilde{Q}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}^\dagger \tilde{U}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}^\dagger \tilde{D}} \\ \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{D}^\dagger \tilde{Q}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{D}^\dagger \tilde{U}} & \left(\widetilde{\mathbf{M}}^2\right)_{\tilde{D}^\dagger \tilde{D}} \end{pmatrix}_{ij} , \tag{2.4}$$

with

$$\begin{aligned}
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{Q}_i^\dagger \tilde{Q}_j} &= \left(\widetilde{\mathbf{M}}_Q^2\right)_{ij} \mathbf{1}_2 + \left(\mathbf{h}_d^\dagger \mathbf{h}_d\right)_{ij} \Phi_1 \Phi_1^\dagger + \left(\mathbf{h}_u^\dagger \mathbf{h}_u\right)_{ij} \left(\Phi_2^\dagger \Phi_2 \mathbf{1}_2 - \Phi_2 \Phi_2^\dagger\right) \\
&\quad - \frac{1}{2}\delta_{ij} g_2^2 \left(\Phi_1 \Phi_1^\dagger - \Phi_2 \Phi_2^\dagger\right) + \delta_{ij} \left(\frac{1}{4} g_2^2 - \frac{1}{12} {g_1}^2\right) \left(\Phi_1^\dagger \Phi_1 - \Phi_2^\dagger \Phi_2\right) \mathbf{1}_2, \\
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}_i^\dagger \tilde{Q}_j} &= \left(\widetilde{\mathbf{M}}^2\right)^\dagger_{\tilde{Q}_j^\dagger \tilde{U}_i} = -\left(\mathbf{a}_u\right)_{ij} \Phi_2^T i\tau_2 + \left(\mathbf{h}_u\right)_{ij} \lambda S^* \Phi_1^T i\tau_2, \\
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{D}_i^\dagger \tilde{Q}_j} &= \left(\widetilde{\mathbf{M}}^2\right)^\dagger_{\tilde{Q}_j^\dagger \tilde{D}_i} = +\left(\mathbf{a}_d\right)_{ij} \Phi_1^\dagger - \left(\mathbf{h}_d\right)_{ij} \lambda S^* \Phi_2^\dagger, \\
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}_i^\dagger \tilde{U}_j} &= \left(\widetilde{\mathbf{M}}_U^2\right)_{ij} + \left(\mathbf{h}_u \mathbf{h}_u^\dagger\right)_{ij} \Phi_2^\dagger \Phi_2 + \frac{1}{3}\delta_{ij} {g_1}^2 \left(\Phi_1^\dagger \Phi_1 - \Phi_2^\dagger \Phi_2\right), \\
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{D}_i^\dagger \tilde{D}_j} &= \left(\widetilde{\mathbf{M}}_D^2\right)_{ij} + \left(\mathbf{h}_d \mathbf{h}_d^\dagger\right)_{ij} \Phi_1^\dagger \Phi_1 - \frac{1}{6}\delta_{ij} {g_1}^2 \left(\Phi_1^\dagger \Phi_1 - \Phi_2^\dagger \Phi_2\right), \\
\left(\widetilde{\mathbf{M}}^2\right)_{\tilde{U}_i^\dagger \tilde{D}_j} &= \left(\widetilde{\mathbf{M}}^2\right)^\dagger_{\tilde{D}_j^\dagger \tilde{U}_i} = \left(\mathbf{h}_u \mathbf{h}_d^\dagger\right)_{ij} \Phi_1^T i\tau \Phi_2 \ .
\end{aligned} \tag{2.5}$$

In the above, $\mathbf{h}_u$ is the 3×3 up-type Yukawa coupling matrix, $\widetilde{\mathbf{M}}^2_{Q,U,D}$ are the 3×3 soft mass-squared matrices of the squarks and $\mathbf{a}_{d,u} = \mathbf{h}_{d,u} \mathbf{A}_{d,u}$ are the corresponding 3×3 soft Yukawa matrices. Notice that (2.4) reduces to the MSSM result presented in [7], after identifying the effective μ-parameter as $\mu = -\frac{1}{\sqrt{2}} \lambda v_S$, where $v_S = \sqrt{2} \langle S \rangle$ is the VEV of S.

The weak left- and right-handed quarks, denoted as $u^0_{L,R}$, $d^0_{L,R}$, are related to the respective mass eigenstates $u_{L,R}$ and $d_{L,R}$ by the unitary transformations

$$u^0_L = \mathbf{U}^Q_L\ u_L, \quad d^0_L = \mathbf{U}^Q_L\ \mathbf{V}\ d_L, \quad u^0_R = \mathbf{U}^u_R\ u_R, \quad d^0_R = \mathbf{U}^d_R\ d_R. \tag{2.6}$$

Here $\mathbf{U}^Q_L$ and $\mathbf{U}^{u,d}_R$ are unitary 3×3 matrices, and $\mathbf{V}$ is the physical CKM-mixing matrix. The matrices $\mathbf{U}^Q_L$ and $\mathbf{U}^{u,d}_R$ are determined by the mass renormalisation conditions

$$\left\langle \mathcal{L}^d_{\text{eff}}\left[\Phi_1,\Phi_2,S\right]\right\rangle \ =\ -\bar{d}_R\widehat{\mathbf{M}}_d d_L\ +\ \text{H.c.}, \quad \left\langle \mathcal{L}^u_{\text{eff}}\left[\Phi_1,\Phi_2,S\right]\right\rangle \ =\ -\bar{u}_R\widehat{\mathbf{M}}_u u_L\ +\ \text{H.c.}, \tag{2.7}$$

where $\langle\ldots\rangle$ indicates the vacuum expectation value of the enclosed expression. In (2.7), $\mathbf{M}_{d,u}$ are the physical down- and up-quark mass matrices and $\mathcal{L}^u_{\text{eff}}$ is the corresponding effective Lagrangian for up-type Yukawa sector [cf. (2.1)]. Imposing the conditions (2.7) yields

$$\mathbf{U}^{d\dagger}_R\ \mathbf{h}_d\ \mathbf{U}^Q_L \ =\ \frac{\sqrt{2}}{v_1}\widehat{\mathbf{M}}_d\ \mathbf{V}^\dagger\ \mathbf{R}_d^{-1}, \quad \mathbf{U}^{u\dagger}_R\ \mathbf{h}_u\ \mathbf{U}^Q_L \ =\ \frac{\sqrt{2}}{v_2}\widehat{\mathbf{M}}_u\ \mathbf{R}_u^{-1}, \tag{2.8}$$

where

$$\begin{aligned}
\mathbf{R}_d \ &=\ \mathbf{1}_3\ +\ \frac{\sqrt{2}}{v_1}\mathbf{U}^{Q\dagger}_L\left\langle \mathbf{h}_d^{-1}\Delta\mathbf{h}_d\left[\Phi_1,\Phi_2,S\right]\right\rangle \mathbf{U}^Q_L\ ,\\
\mathbf{R}_u \ &=\ \mathbf{1}_3\ +\ \frac{\sqrt{2}}{v_2}\mathbf{U}^{Q\dagger}_L\left\langle \mathbf{h}_u^{-1}\Delta\mathbf{h}_u\left[\Phi_1,\Phi_2,S\right]\right\rangle \mathbf{U}^Q_L\ .
\end{aligned} \tag{2.9}$$

Note that $\Delta\mathbf{h}_u$ is the respective 1-loop effective functional for the up-type quark Yukawa couplings.

Considering now the general effective FCNC Lagrangian (2.1) for Higgs interactions to down-type quarks, we find that

$$\begin{aligned}
-\mathcal{L}^{d,H}_{\text{FCNC}} \ &=\ \bar{d}_R\frac{\mathbf{h}_d}{\sqrt{2}}\Big[\phi_1\left(\mathbf{1}_3+\mathbf{\Delta}_d^{\phi_1}\right)-ia_1\left(\mathbf{1}_3+\mathbf{\Delta}_d^{a_1}\right)\\
&\qquad +\phi_2\mathbf{\Delta}_d^{\phi_2}-ia_2\mathbf{\Delta}_d^{a_2}+\phi_S\mathbf{\Delta}_d^{\phi_S}-ia_S\mathbf{\Delta}_d^{a_S}\Big]\mathbf{V}d_L\\
&\qquad +\bar{d}_R\mathbf{h}_d\left[\phi_1^-\left(\mathbf{1}_3+\mathbf{\Delta}_d^{\phi_1^-}\right)+\phi_2^-\mathbf{\Delta}_d^{\phi_2^-}\right]u_L\ +\ \text{H.c.},
\end{aligned} \tag{2.10}$$

where the component Higgs fields are given by

$$\Phi_{1,2} \ =\ \begin{pmatrix} \phi^+_{1,2} \\ \frac{1}{\sqrt{2}}\left(v_{1,2}+\phi_{1,2}+ia_{1,2}\right)\end{pmatrix}, \qquad S\ =\ \frac{1}{\sqrt{2}}\left(v_S+\phi_S+ia_S\right). \tag{2.11}$$

In (2.10), $\mathbf{\Delta}_d^{\phi_{1,2,S}}$, $\mathbf{\Delta}_d^{a_{1,2,S}}$ and $\mathbf{\Delta}_d^{\phi^\pm_{1,2}}$ are 3-by-3 matrices that are evaluated as

$$\mathbf{\Delta}_d^{\phi_{1,2,S}} = \sqrt{2}\left\langle\frac{\delta}{\delta\phi_{1,2,S}}\mathbf{\Delta}_d\right\rangle, \quad \mathbf{\Delta}_d^{a_{1,2,S}} = i\sqrt{2}\left\langle\frac{\delta}{\delta a_{1,2,S}}\mathbf{\Delta}_d\right\rangle, \quad \mathbf{\Delta}_d^{\phi^\pm_{1,2}} = \left\langle\frac{\delta}{\delta\phi^\pm_{1,2}}\mathbf{\Delta}_d\right\rangle, \tag{2.12}$$

where we have defined $\mathbf{\Delta}_d \equiv \mathbf{h}_d^{-1}\Delta\mathbf{h}_d[\Phi_1,\Phi_2,S]$ and suppressed the vanishing iso-doublet components on the LHS of (2.12). Since the singlet Higgs field does not contain a charged

component, the couplings of the charged Higgs boson are unchanged from those of the MSSM and we shall not consider them here any further. In the neutral sector, the physical Higgs boson mass eigenstates are related to the weak eigenstates by the orthogonal mixing matrices O^H and O^A, such that

$$\begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_S \end{pmatrix} = O^H \begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix}, \qquad \begin{pmatrix} a \\ a_S \end{pmatrix} = O^A \begin{pmatrix} A_1 \\ A_2 \end{pmatrix} . \tag{2.13}$$

It is sometimes useful to parameterise O^A by the CP-odd mixing angle θ_A, where

$$O^A = \begin{pmatrix} \cos\theta_A & \sin\theta_A \\ -\sin\theta_A & \cos\theta_A \end{pmatrix} . \tag{2.14}$$

In terms of the mass eigenstates, the general effective Lagrangian (2.1) that includes the FCNC interactions of the neutral Higgs bosons with the down-type quarks reads:

$$\begin{aligned} -\mathcal{L}^{d,H}_{\rm FCNC} &= \frac{g}{2M_W}\Big[H_i\bar{d}_R\left(\widehat{\mathbf{M}}_d\ \mathbf{g}^L_{H_i\bar{d}d}P_L + \mathbf{g}^R_{H_i\bar{d}d}\ \widehat{\mathbf{M}}_d P_R\right) d \\ &\quad + A_j\bar{d}_R\left(\widehat{\mathbf{M}}_d\ \mathbf{g}^L_{A_j\bar{d}d}P_L + \mathbf{g}^R_{A_j\bar{d}d}\ \widehat{\mathbf{M}}_d P_R\right) d\Big] , \end{aligned} \tag{2.15}$$

where the Higgs couplings in the flavour basis $\mathbf{U}^Q_L = \mathbf{U}^u_R = \mathbf{U}^d_R = \mathbf{1}_3$ are given by

$$\begin{aligned} \mathbf{g}^L_{H_i\bar{d}d} &= \frac{\mathcal{O}^H_{1i}}{c_\beta}\mathbf{V}^\dagger\mathbf{R}_d^{-1}\left(\mathbf{1}_3 + \mathbf{\Delta}_d^{\phi_1}\right)\mathbf{V} + \frac{\mathcal{O}^H_{2i}}{c_\beta}\mathbf{V}^\dagger\mathbf{R}_d^{-1}\mathbf{\Delta}_d^{\phi_2}\mathbf{V} \\ &\quad + \frac{\mathcal{O}^H_{3i}}{c_\beta}\mathbf{V}^\dagger\mathbf{R}_d^{-1}\mathbf{\Delta}_d^{\phi_S}\mathbf{V} , \end{aligned} \tag{2.16}$$

$$\mathbf{g}^R_{H_i\bar{d}d} = \left(\mathbf{g}^L_{H_i\bar{d}d}\right)^\dagger , \tag{2.17}$$

$$\begin{aligned} \mathbf{g}^L_{A_i\bar{d}d} &= i\mathcal{O}^A_{1i}t_\beta\mathbf{V}^\dagger\mathbf{R}_d^{-1}\left(\mathbf{1}_3 + \mathbf{\Delta}_d^{a_1} - \frac{1}{t_\beta}\mathbf{\Delta}_d^{a_2}\right)\mathbf{V} \\ &\quad - i\frac{\mathcal{O}^A_{2i}}{c_\beta}\mathbf{V}^\dagger\mathbf{R}_d^{-1}\mathbf{\Delta}_d^{a_S}\mathbf{V} , \end{aligned} \tag{2.18}$$

$$\mathbf{g}^R_{A_i\bar{d}d} = \left(\mathbf{g}^L_{A_i\bar{d}d}\right)^\dagger . \tag{2.19}$$

In the above, we used the short-hand notation: $t_\beta \equiv \tan\beta = v_2/v_1$, $c_\beta \equiv \cos\beta$ and $s_\beta \equiv \sin\beta$.

2.1 Single Higgs Insertion Approximation

Our numerical results of Section 4 include a full evaluation of $\mathbf{\Delta}_d$ and its derivatives using the matrix expression of (2.2). We may obtain an understanding of our results by considering an expansion of $\Delta\mathbf{h}_d$ in terms of Φ_2 and S. We call the leading term of such an expansion, as represented by Fig. 1, the single-Higgs-insertion (SHI) approximation.

Within the SHI approximation, the $\tan\beta$-enhanced threshold corrections satisfy the simple relation

$$\frac{\sqrt{2}}{v_2}\langle\mathbf{\Delta}_d\rangle = \mathbf{\Delta}_d^{\phi_2} = \mathbf{\Delta}_d^{a_2} = \frac{v_S}{v_2}\mathbf{\Delta}_d^{\phi_S} = \frac{v_S}{v_2}\mathbf{\Delta}_d^{a_S} \ , \tag{2.20}$$

where

$$\frac{\sqrt{2}}{v_2}\langle\mathbf{\Delta}_d\rangle \ = \ \mathbf{1}_3\frac{2\alpha_S}{3\pi}\mu M_3 I(\widetilde{M}_Q^2,\widetilde{M}_D^2,M_3^2) \ + \ \frac{\mathbf{h}_u^\dagger\mathbf{h}_u}{16\pi^2}\mu A_u I(\widetilde{M}_Q^2,\widetilde{M}_U^2,\mu^2) \ , \tag{2.21}$$

and $I(x,y,z)$ is the 1-loop function,

$$I(x,y,z) = \frac{xy\ln(x/y)+yz\ln(y/z)+xz\ln(z/x)}{(x-y)(y-z)(x-z)} \ . \tag{2.22}$$

In writing the above, we have neglected the subdominant contributions coming from chargino- and neutralino-exchange graphs, which are weak-coupling constant suppressed.

3 B-meson FCNC Observables

In this section we review the analytic results relevant to FCNC B-meson observables. Our conventions and discussion follow [6].

3.1 ΔM_{B_q}

In the approximation of equal B-meson lifetimes, the SM and SUSY contributions may be written separately as

$$\Delta M_{B_q} = 2\left|\left\langle\bar{B}_q^0\right|H_{\rm eff}^{\Delta B=2}\left|B_q^0\right\rangle_{\rm SM} + \left\langle\bar{B}_q^0\right|H_{\rm eff}^{\Delta B=2}\left|B_q^0\right\rangle_{\rm SUSY}\right| \ . \tag{3.1}$$

Although the SM predictions are consistent with the observed experimental values [22], the uncertainties are large and a non-negligible SUSY contribution is not excluded.

The SUSY contributions are given by

$$\begin{aligned}
\left\langle \bar{B}_d^0 \right| H_{\rm eff}^{\Delta B=2} \left| B_d^0 \right\rangle_{\rm SUSY} &= 1711 {\rm ps}^{-1} \left(\frac{\hat{B}_{B_d}^{1/2} F_{B_d}}{230 \text{ MeV}} \right)^2 \left(\frac{\eta_B}{0.55} \right) \\
&\times \left[0.88 \left(C_2^{\rm LR(DP)} + C_2^{\rm LR(2HDM)} \right) - 0.52 \left(C_1^{\rm SLL(DP)} + C_1^{\rm SRR(DP)} \right) \right] \ , \quad (3.2) \\
\left\langle \bar{B}_s^0 \right| H_{\rm eff}^{\Delta B=2} \left| B_s^0 \right\rangle_{\rm SUSY} &= 2310 {\rm ps}^{-1} \left(\frac{\hat{B}_{B_s}^{1/2} F_{B_s}}{265 \text{ MeV}} \right)^2 \left(\frac{\eta_B}{0.55} \right) \\
&\times \left[0.88 \left(C_2^{\rm LR(DP)} + C_2^{\rm LR(2HDM)} \right) - 0.52 \left(C_1^{\rm SLL(DP)} + C_1^{\rm SRR(DP)} \right) \right] \ , \quad (3.3)
\end{aligned}$$

where DP stands for the Higgs-mediated double-penguin contributions. Here F_{B_q} is the weak decay constant and B_{B_q} is the so-called "Bag" parameter of the B_q^0 meson respectively. The numerical factor η_B is due to QCD corrections. We use the next-to-leading order QCD factors [5, 23] and hadronic matrix elements at the scale $\mu = 4.2$ GeV:

$$\bar{P}_1^{\rm LR} = -0.58 \ , \qquad \bar{P}_2^{\rm LR} = 0.88 \ , \qquad \bar{P}_1^{\rm SLL} = -0.52 \ , \qquad \bar{P}_2^{\rm SLL} = -1.1 \ . \tag{3.4}$$

The double-penguin Wilson coefficients in (3.2) are given by

$$\begin{aligned}
C_1^{\rm SLL(DP)} &= \frac{-16\pi^2 m_b^2}{\sqrt{2} G_F M_W^2} \left(\sum_{i=1}^{3} \frac{\mathbf{g}^L_{H_i \bar{b} q} \mathbf{g}^L_{H_i \bar{b} q}}{M_{H_i}^2} + \sum_{j=1}^{2} \frac{\mathbf{g}^L_{A_j \bar{b} q} \mathbf{g}^L_{A_j \bar{b} q}}{M_{A_j}^2} \right) , \\
C_1^{\rm SRR(DP)} &= \frac{-16\pi^2 m_q^2}{\sqrt{2} G_F M_W^2} \left(\sum_{i=1}^{3} \frac{\mathbf{g}^R_{H_i \bar{b} q} \mathbf{g}^R_{H_i \bar{b} q}}{M_{H_i}^2} + \sum_{j=1}^{2} \frac{\mathbf{g}^R_{A_j \bar{b} q} \mathbf{g}^R_{A_j \bar{b} q}}{M_{A_j}^2} \right) , \\
C_2^{\rm LR(DP)} &= \frac{-32\pi^2 m_b m_q}{\sqrt{2} G_F M_W^2} \left(\sum_{i=1}^{3} \frac{\mathbf{g}^L_{H_i \bar{b} q} \mathbf{g}^R_{H_i \bar{b} q}}{M_{H_i}^2} + \sum_{j=1}^{2} \frac{\mathbf{g}^L_{A_j \bar{b} q} \mathbf{g}^R_{A_j \bar{b} q}}{M_{A_j}^2} \right) ,
\end{aligned} \tag{3.5}$$

where the relevant couplings $\mathbf{g}^{L,R}_{H_i(A_j)\bar{b}q}$ are given in (2.16). We neglect the B-meson masses and Higgs boson widths in the denominators of the sums, except for the lightest Higgs mass eigenstates H_1 and A_1, since we may have masses $\mathcal{O}(1 \text{ GeV})$ in the case of a highly gauge singlet particle. In this limit we replace the effective propagators $M^{-2}_{H_1(A_1)}$ by their Breit-Wigner forms. Of the two relevant 1-loop contributions to $\left\langle \bar{B}^0 | H_{\rm eff}^{\Delta B=2} | B^0 \right\rangle$, the $t - H^\pm$ box contribution to $C_2^{\rm LR}$ may be given to a good approximation by [5]

$$C_2^{\rm LR(2HDM)} \approx -\frac{2 m_b m_q}{M_W^2} (V_{tb}^* V_{tq})^2 \tan^2 \beta \ . \tag{3.6}$$

We also include the chargino-stop box diagram which contributes to $C_1^{\rm SLL}$, although its contribution remains subdominant.

3.2 $\bar{B}^0_{d,s} \to \mu^+\mu^-$

Neglecting contributions proportional to the lighter quark masses $m_{d,s}$, the effective Hamiltonian for $\Delta B = 1$ FCNC processes is given by

$$H_{\rm eff}^{\Delta B=1} \;=\; -2\sqrt{2}G_F V_{tb}V_{tq}^* \left(C_S\mathcal{O}_S + C_P\mathcal{O}_P + C_{10}\mathcal{O}_{10}\right) \;, \tag{3.7}$$

where

$$\begin{aligned}
\mathcal{O}_S &= \frac{e^2}{16\pi^2} m_b \left(\bar{q}P_R b\right)\left(\bar{\mu}\mu\right) \;, \\
\mathcal{O}_P &= \frac{e^2}{16\pi^2} m_b \left(\bar{q}P_R b\right)\left(\bar{\mu}\gamma_5\mu\right) \;, \\
\mathcal{O}_{10} &= \frac{e^2}{16\pi^2} m_b \left(\bar{q}\gamma^\mu P_L b\right)\left(\bar{\mu}\gamma_\mu\gamma_5\mu\right) \;.
\end{aligned} \tag{3.8}$$

The Wilson coefficients C_S and C_P are given at large $\tan\beta$ by

$$\begin{aligned}
C_S &= \frac{2\pi m_\mu}{\alpha_{\rm em}} \frac{1}{V_{tb}V_{tq}^*} \sum_{i=1}^{3} \frac{\mathbf{g}^R_{H_i\bar{q}b}\mathbf{g}^S_{H_i\bar{\mu}\mu}}{M^2_{H_i}} \;, \\
C_P &= i\,\frac{2\pi m_\mu}{\alpha_{\rm em}} \frac{1}{V_{tb}V_{tq}^*} \sum_{i=1}^{2} \frac{\mathbf{g}^R_{A_i\bar{q}b}\mathbf{g}^P_{A_i\bar{\mu}\mu}}{M^2_{A_i}} \;,
\end{aligned} \tag{3.9}$$

and $C_{10} = -4.221$ denotes the leading SM contribution. Again we neglect the B-meson masses and Higgs boson widths except for the contribution of the lightest Higgs scalar and pseudoscalar. The reduced scalar and pseudoscalar Higgs couplings to charged leptons $\mathbf{g}^{S(P)}_{H_i(A_i)\bar{l}l}$ are given in [18]. The leptonic vertex corrections can become important in the limit of a light, highly singlet Higgs boson, so we retain them in our numerical estimates.

Using the effective $\Delta B = 1$ Hamiltonian (3.7), the branching ratio for $\bar{B}^0_{d,s} \to \mu^+\mu^-$ is given by [24]

$$\mathcal{B}(\bar{B}^0_{d,s} \to \mu^+\mu^-) = \frac{G_F^2\alpha_{\rm em}^2}{16\pi^3} M_{B_q}\tau_{B_q}|V_{tb}V_{tq}^*|^2 \sqrt{1-\frac{4m_\mu^2}{M^2_{B_q}}} \left[\left(1-\frac{4m_\mu^2}{M^2_{B_q}}\right)|F_S^q|^2 + |F_P^q + 2m_\mu F_A^q|^2\right], \tag{3.10}$$

where $q = d, s$ and τ_{B_q} is the total lifetime of the B_q meson. The form factors $F^q_{S,P,A}$ are given by

$$F^q_{S,P} = -\frac{i}{2} M^2_{B_q} F_{B_q} \frac{m_b}{m_b + m_q} C_{S,P} \;, \qquad F^q_A = -\frac{i}{2} F_{B_q} C_{10} \;. \tag{3.11}$$

4 Numerical Results

In this section we provide numerical estimates of FCNC observables, within the contexts of both the MNSSM and NMSSM. We focus on those sectors of the models where the gauge-singlet Higgs fields are predicted to be light. We assume the framework of MFV so that all flavour changing effects are proportional to the CKM matrix $\mathbf{V}$. This implies that the quark-sector soft SUSY-breaking terms are proportional to $\mathbf{1}_3$ [7], i.e.

$$\widetilde{\mathbf{M}}^2_{Q,U,D} = \widetilde{M}^2_{Q,U,D}\ \mathbf{1}_3, \qquad \mathbf{A}_{u,d} = A_{u,d}\ \mathbf{1}_3\ , \tag{4.1}$$

and similarly for the leptonic sector. We neglect the neutralino-mediated contributions to the FCNC Higgs boson couplings, which are found to be subdominant.

Important constraints on FCNCs come from $B \to X_s\gamma$ experiments, which are sensitive to large stop-quark mixing due to a dominant chargino-stop quark loop. To avoid conflict with these constraints, we assume that $M_{\rm SUSY}$ is relatively heavy of order ~ 2 TeV. In calculating the Higgs couplings, we have used the following benchmark values for the soft SUSY-breaking parameters

$$\begin{aligned} &\widetilde{M}^2_Q = \widetilde{M}^2_L = \widetilde{M}^2_D = \widetilde{M}^2_E = \widetilde{M}^2_U \ = (1.7\ \text{TeV})^2, \\ &A_u = A_d = A_e = 2.0\ \text{TeV}, \\ &M_1 = M_2 = M_3 = 2.0\ \text{TeV}\ . \end{aligned} \tag{4.2}$$

In addition, we take $\mu = 140$ GeV and $t_\beta = 50$ throughout. The Higgs-mediated contribution to $B \to X_s\gamma$ proceeds through the charged Higgs boson and the predictions of the MSSM are not altered by the presence of the singlet Higgs bosons. We have made use of the public code CPsuperH [25] to check that all points considered here are consistent with the 2σ experimental bounds on $B \to X_s\gamma$; the Higgs-mediated contribution was found to be subdominant in the region of interest.

4.1 Electroweak-scale Higgs singlets in the MNSSM

The renormalisable MNSSM superpotential is given by

$$\mathcal{W}_{\rm MNSSM} = \mathcal{W}_{\rm Yuk} + \lambda \hat{S}\hat{H}_1\hat{H}_2 + t_F\hat{S}\ , \tag{4.3}$$

where $\mathcal{W}_{\rm Yuk}$ represents the Yukawa couplings of the MSSM superpotential. The tadpole parameter t_F and its associated soft SUSY-breaking term t_S are radiatively generated by supergravity quantum effects from Planck-suppressed non-renormalisable operators in the Kähler potential and superpotential. Discrete R-symmetries, such as Z_5^R and Z_7^R, that are imposed on the theory postpones the appearance of these operators beyond the 5-loop level, such that they are naturally suppressed of the order of $M_{\rm SUSY}$ within a perturbative

framework of supergravity (SUGRA). For a detailed proof of the argument, see the first reference of [11].

The tree-level Higgs sector of the MNSSM may be described by the six parameters

$$t_\beta,\ M_a,\ \mu,\ \lambda,\ m_{12}^2,\ \frac{\lambda t_S}{\mu}\ , \tag{4.4}$$

where M_a is the would-be MSSM pseudoscalar Higgs mass and $m_{12}^2 = \lambda t_F$. In our numerical results we also include the dominant radiative corrections due to both (s)top and (s)bottom loops, which are necessary to raise the mass of the SM-like Higgs boson above the LEP II constraint of 114 GeV.

Our interest here is in the effects of light gauge-singlet Higgs bosons, with masses at or below the electroweak scale. Such a scenario may be realised by taking M_a to be large, so that the heavy Higgs doublet fields effectively decouple, and by choosing m_{12}^2 such that the mixing between the light CP-even Higgs bosons is suppressed. In the numerical estimates presented here, we take

$$M_a = 1.5 \text{ TeV}, \quad m_{12}^2 = (1.0 \text{ TeV})^2, \quad \mu = 140 \text{ GeV}, \quad \tan\beta = 50, \quad \lambda = 0.3\ . \tag{4.5}$$

This corresponds to a Higgs pseudoscalar mixing angle of $\cos\theta_A \sim 0.17$. The scalar Higgs singlet mixes almost exclusively with the ϕ_1 doublet, the non-singlet fraction is approximately equal for both the lightest Higgs states H_1 and A_1. These light particles are not constrained by the direct search limits from LEP II and the Tevatron due to their highly gauge-singlet nature, with $g_{H_{1,2}ZZ},\ g_{H_1A_1Z} < 0.05$. The mass scale of the singlet Higgs bosons are approximately given by $\sqrt{\lambda t_S/\mu}$. Notice that, in the absence of strong singlet-doublet mixing effects, the singlet Higgs scalar and pseudoscalar are approximately degenerate due to the tree-level mass-sum rule [11]:

$$m_{H_1}^2 \ + \ m_{H_2}^2 \ + \ m_{H_3}^2 \ = \ M_Z^2 \ + \ m_{A_1}^2 \ + \ m_{A_2}^2\ . \tag{4.6}$$

Including dominant radiative corrections we find the masses of the remaining Higgs bosons in this scenario to be

$$m_{H_2} = 147 \text{ GeV}, \quad m_{H_3} = 1.52 \text{ TeV}, \quad m_{A_2} = 1.52 \text{ TeV}, \tag{4.7}$$

so that the H_2 Higgs boson, which has SM-like couplings to the EW gauge bosons, is well above the direct search limit set by LEP II.

4.1.1 Effects on ΔM_{B_q}

In Fig. 2 we show the SUSY contribution to ΔM_{B_s} in units of ps^{-1}, as a function of the lightest pseudoscalar mass. The upper curve fully includes the radiative Yukawa couplings




Figure 2: *The SUSY contribution to* ΔM_{B_s} *in units of* ps^{-1} *as a function of the mass of the lightest pseudoscalar* m_{A_1} *in the MNSSM. All parameters are taken as in (4.2) and (4.5). The horizontal lines show the currently measured value along with the SM and MSSM predictions for corresponding values of* M_a *and* $\tan\beta$.

of the Higgs singlet fields whilst the lower curve neglects these corrections. We observe that the SUSY contribution can be larger than the currently observed value at large $\tan\beta$ if the singlet Higgs bosons are light. For moderate values of m_{A_1}, the effects of the direct singlet Yukawa coupling remain significant, enhancing the value of ΔM_{B_s} by around 25%.

The dominant contribution to ΔM_{B_s} for light singlet Higgs bosons is due to the Wilson coefficient $C_1^{\mathrm{SLL(DP)}}$. Considering the SHI approximation for the reduced couplings $\mathbf{g}^L_{H_1(A_1)\bar{b}q}$, we see that the off-diagonal terms are given by

$$\mathbf{g}^L_{H_1\bar{b}q} = \left(t_\beta \mathcal{O}^H_{11} - \frac{v}{v_S}\mathcal{O}^H_{31}\right)\mathbf{V}^\dagger\mathbf{R}^{-1}\mathbf{V}\ , \tag{4.8}$$

$$\mathbf{g}^L_{A_1\bar{b}q} = i\left(t_\beta \mathcal{O}^A_{11} + \frac{v}{v_S}\mathcal{O}^A_{21}\right)\mathbf{V}^\dagger\mathbf{R}^{-1}\mathbf{V}\ , \tag{4.9}$$

where we have neglected the small contributions of the Φ_2 component next to those of Φ_1.




Figure 3: *The dominant Wilson coefficients contributing to ΔM_{B_s} as a function of the mass of the lightest pseudoscalar m_{A_1} in the MNSSM. The dashed curve shows the Wilson coefficient $C_1^{\rm SLL(DP)}$ neglecting the threshold corrections for the gauge singlet Higgs boson.*

We observe that the FCNC couplings of the lightest scalar and pseudoscalar are of equal magnitude, if threshold corrections to the singlet-Higgs Yukawa couplings are neglected and $\mathcal{O}_{11}^H \simeq \mathcal{O}_{11}^A$. In this case, there is a cancellation between the H_1- and the A_1-mediated contributions to $C_1^{\rm SLL(DP)}$. An analogous cancellation between the heavy Higgs bosons H and A is known to take place in the MSSM as well, as a consequence of the PQ symmetry which forbids at the tree-level the dominant $\Delta B = 2$ operators, $(\bar{b}_R d_L)^2$ and $(\bar{b}_R s_L)^2$, associated with $C_1^{\rm SLL(DP)}$ (see, e.g. [5]).

For very light singlets this cancellation is dominantly broken by the mass splitting between H_1 and A_1, typically of the order $m_{H_1} \sim m_{A_1} + 2$ GeV. For larger masses, this splitting is negligible and the dominant breaking is instead due to threshold effects on the singlet-Higgs Yukawa couplings, which are found to contribute to the reduced couplings with opposite sign, due to the mixing matrices. The dominant Wilson coefficients for ΔM_{B_s} are plotted in Fig. 3. We do not plot $C_1^{\rm SRR(DP)}$, which exhibits a behaviour similar




Figure 4: *The SUSY contribution to* ΔM_{B_d} *in units of* ps^{-1} *as a function of the mass of the lightest pseudoscalar* m_{A_1} *in the MNSSM. The line conventions are as for Fig. 2. All parameters are taken as in (4.2) and (4.5). The upper horizontal line shows the currently measured value and the lower horizontal line shows the MSSM prediction for corresponding values of* M_a *and* $\tan\beta$*. We do not show the SM prediction as the central value is close to the experimentally observed splitting.*

to $C_1^{\mathrm{SLL(DP)}}$, but which is suppressed by a factor of $(m_s/m_b)^2$.

In Fig. 4 we show the corresponding plot for the contribution to ΔM_{B_d}. Again we observe that the SUSY contribution can exceed the currently observed value, with the limit on m_{A_1} at the same level as that due to ΔM_{B_s}. The effects of the singlet Yukawa coupling are more pronounced at larger values of m_{A_1} than in the case of ΔM_{B_s}, as the enhanced Wilson coefficient $C_1^{\mathrm{SLL(DP)}}$ remains dominant over the two-Higgs-doublet model contribution $C_2^{\mathrm{LR(2HDM)}}$ by an order of magnitude here even for $m_{A_1} \sim 100$ GeV.




Figure 5: *The branching ratio $\mathcal{B}(B_s \to \mu^+\mu^-)$ as a function of the mass of the lightest pseudoscalar m_{A_1} in the MNSSM. All parameters are taken as in (4.2) and (4.5). The horizontal lines show the SM [26] and MSSM predictions and the current experimental upper limit at 90% C.L.*

4.1.2 Effects on $\bar{B}^0_{d,s} \to \mu^+\mu^-$

Figure 5 shows the branching ratio $\mathcal{B}(B_s \to \mu^+\mu^-)$, as a function of m_{A_1}. As expected, the prediction varies strongly with the mass of the light gauge singlet Higgs bosons due to the $M^{-4}_{H_1(A_1)}$ dependence of the form factors $F^s_{S(P)}$. The prediction exceeds the current bounds for Higgs singlet masses below around 50 GeV. Since the branching ratio in (3.10) depends only on the absolute values of the form factors, there is no cancellation between the scalar and pseudoscalar contributions. At low values of m_{A_1}, including the threshold corrections leads to a slight increase in C_P, which is compensated for by a slight decrease in C_S and the overall effect is negligible. At higher masses, $m_{A_1} \gtrsim 60$ GeV, there is cancellation between the contributions due to C_P and C_{10}. In this region the singlet threshold corrections produce a noticeable shift.

4.2 Light singlet Higgs pseudoscalar in the NMSSM

The NMSSM superpotential is given by

$$\mathcal{W}_{\rm NMSSM} = \mathcal{W}_{\rm Yuk} + \lambda \hat{S}\hat{H}_1\hat{H}_2 + \frac{\kappa}{3}\hat{S}^3 \,. \tag{4.10}$$

The corresponding soft SUSY-breaking terms are given by

$$\mathcal{L}^{\rm soft}_{\rm NMSSM} = \lambda A_\lambda S\Phi_1\Phi_2 + \kappa A_\kappa S^3 \,. \tag{4.11}$$

In the limit $A_{\lambda,\kappa} \to 0$, the scalar potential of the NMSSM possesses an additional $U(1)$ symmetry beyond $U(1)_{\rm PQ}$. This symmetry results from an R-symmetry, denoted by $U(1)_{\rm R}$, which is exact in the absence of soft SUSY-breaking trilinear couplings. Recall that the superpotential of the NMSSM is charged under $\mathrm{U}(1)_{\rm R}$ and so it differs from $U(1)_{\rm PQ}$. This $U(1)_{\rm R}$ is spontaneously broken when S acquires a VEV, so that for small values of $A_{\lambda,\kappa}$ the singlet Higgs pseudoscalar is again a pseudo-Goldstone boson and is naturally expected to be much lighter than the other Higgs fields [11, 27].

In order to examine FCNC observables within such a scenario, we take the CP-odd mixing angle θ_A and the mass of the lightest Higgs pseudoscalar m_{A_1} to be free parameters in place of $A_{\lambda,\kappa}$, along with the couplings λ and κ. We use the following benchmark values throughout this section;

$$\lambda = 0.4\,, \quad \kappa = -0.5\,, \quad \cos\theta_A = 0.018\,, \quad \mu = 140\ \text{GeV}\,, \quad \tan\beta = 50\,, \tag{4.12}$$

which are found to lead to the following masses for the remaining Higgs bosons,

$$m_{H_1} = 135\ \text{GeV}, \quad m_{H_2} = 357\ \text{GeV}, \quad m_{H_3} = 1.14\ \text{TeV}, \quad m_{A_2} = 1.14\ \text{TeV}. \tag{4.13}$$

The lightest CP-even Higgs in this scenario has SM-like couplings to the EW gauge bosons.

For general values of κ, the CP-odd and CP-even singlets of the NMSSM are not constrained to have degenerate masses. In the scenario considered here, there is only one light Higgs particle, the pseudo-Goldstone boson A_1. Due to this, there is no cancellation between the dominant Higgs field contributions to the Wilson coefficients as described in Section 4.1 for the MNSSM. This forces us to take a smaller singlet-doublet mixing parameter $\cos\theta_A$ in order to find phenomenologically acceptable results at large values of $\tan\beta$. Note that the value of $\cos\theta_A$ used here is only $\sim 10\%$ of that found for the MNSSM scenario considered previously and that the contributions to $\mathbf{g}^{L(R)}_{A_1\bar{b}q}$ from singlet-doublet mixing and direct threshold corrections are therefore comparable in magnitude.

4.2.1 Effects on ΔM_{B_q}

In Fig. 6 we show the SUSY contribution to ΔM_{B_s} as a function of the lightest Higgs pseudoscalar mass. In this scenario the SUSY contribution also exceeds the currently measured value of ΔM_{B_s} for the lightest values of m_{A_1} at large values of $\tan\beta$.




Figure 6: *The SUSY contribution to* ΔM_{B_s} *in units of* ps^{-1} *as a function of the mass of the lightest pseudoscalar* m_{A_1} *in the NMSSM. The line conventions are as for Fig. 2. All parameters are taken as in (4.2) and (4.12). The horizontal lines show the currently measured value along with the SM and MSSM predictions for corresponding values of* M_a *and* $\tan\beta$.

We remark that the contribution to ΔM_{B_s} in this scenario of the NMSSM is rather smaller than in the MNSSM scenario discussed in Section 4.1. The leading contribution in both cases is due to the Wilson coefficient $C_1^{\text{SLL(DP)}}$. As discussed above, in the NMSSM there is no cancellation between dominant CP-odd and CP-even Higgs fields and so we have taken the FCNC Yukawa couplings to be an order of magnitude smaller than those considered in the previous section. As a result of this, the contribution of the singlet pseudoscalar become negligible here for $m_{A_1} \gtrsim 25$ GeV and we recover the MSSM prediction.

Figure 7 shows the corresponding predictions for ΔM_{B_d}. As in the MNSSM scenario considered in Section 4.1 the bounds from $B_d - \bar{B}_d$ mixing are at the same level as those from the B_s mesons, although the relative importance of the singlet threshold corrections at larger masses is greater due to the continuing dominance of $C_1^{\text{SLL(DP)}}$ over $C_2^{\text{LR(DP)}}$. In




Figure 7: *The SUSY contribution to* ΔM_{B_d} *in units of* ps^{-1} *as a function of the mass of the lightest pseudoscalar* m_{A_1} *in the NMSSM. The line conventions are as for Fig. 2. All parameters are taken as in (4.2) and (4.12). The upper horizontal line shows the currently measured value and the lower horizontal line shows the MSSM prediction for corresponding values of* M_a *and* $\tan\beta$*. We do not show the SM prediction as the central value is close to the experimentally observed splitting.*

particular, the contributions of the singlet pseudoscalar are not negligible at the same low masses as for ΔM_{B_s}.

4.2.2 Effects on $\bar{B}_s^0 \to \mu^+\mu^-$

Figure 8 shows the branching ratio $\mathcal{B}(B_s \to \mu^+\mu^-)$ as a function of the lightest pseudoscalar mass in the NMSSM. For small values of m_{A_1} the prediction exceeds the current 90% confidence limits at large $\tan\beta$. For a heavier singlet Higgs pseudoscalar, we observe that inclusion of the singlet threshold corrections leads to a significant reduction in the branching ratio, which becomes suppressed by more than an order of magnitude compared




Figure 8: *The branching ratio $\mathcal{B}(B_s \to \mu^+\mu^-)$ as a function of the mass of the lightest pseudoscalar, m_{A_1} in the NMSSM. The solid line includes threshold corrections for the gauge singlet Higgs bosons, the dashed curve neglects these. All parameters are taken as in (4.2) and (4.12). The dot-dashed curve shows the NMSSM prediction, including all threshold corrections, but with $A_u = -2$ TeV. The horizontal lines show the SM [26] and the MSSM predictions and the current experimental upper limit at 90% C.L. respectively.*

to the SM prediction. In this mass range the contribution of the light pseudoscalar interferes destructively with the SM-like diagrams. It should be noted that such a large suppression is not possible in either the MSSM or MNSSM, since in both these models tree-level mass-sum rules prevent the appearance of isolated CP-odd Higgs bosons.

The leading effects of the neutral Higgs bosons on $\mathcal{B}(B_s \to \mu^+\mu^-)$ in both the MSSM and MNSSM may be understood by including only the contributions due to the dominant scalar/pseudoscalar pair (i.e. H, A in the MSSM or H_1, A_1 in the above considered limit of the MNSSM) and by assuming equal masses and effective Yukawa couplings (this is accurate to the order of a few percent). In this approximation, it can be shown that the absolute minimum of $\mathcal{B}(B_s \to \mu^+\mu^-)$ in both these models is $1/2 \times \mathcal{B}(B_s \to \mu^+\mu^-)_{\rm SM}$.

When such a cancellation is effective in the NMSSM, the absolute minimum of $\mathcal{B}(B_s \to \mu^+\mu^-)$ is due to the contribution of the heavy CP-even Higgs doublet, $H_3 \sim \phi_1$. The exact value of $\mathcal{B}(B_s \to \mu^+\mu^-)$ at the minimum scales as $M_{H_3}^{-4}$ and the branching ratio will be further suppressed as H_3 and A_2 become heavier. As can be seen from the dot-dashed curve in Fig. 8, the suppression is found to vanish when μA_u flips sign, i.e. for $A_u = -2$ TeV, since the leading SM and SUSY contributions interfere constructively in this case.

5 Conclusions

We have shown that singlet Higgs bosons generically present in extensions of the MSSM can contribute significantly to FCNC observables at large $\tan\beta$, both through their tree-level mixing with the Higgs doublet fields and through 1-loop quantum effects. For very light Higgs singlets, the contributions may exceed even by one order of magnitude the current experimental limits on FCNC processes. This allows us to place severe constraints on the mass of the lightest scalar state in the theory at large values of $\tan\beta$.

In the MNSSM, the scalar and pseudoscalar singlet Higgs fields are constrained to be approximately degenerate by a tree-level mass sum rule [11]. Their contributions to B-meson mixing through the dominant Wilson coefficient $C_1^{\rm SLL}$ interfere destructively, analogously to the contributions of the MSSM heavy Higgs bosons. At moderate to large values of the common Higgs singlet mass, the direct coupling plays an important role in lifting this cancellation, leading to a large enhancement of $C_1^{\rm SLL}$. Although cancellations between the pseudoscalar and SM-like contributions to the branching ratio $B_s \to \mu^+\mu^-$ are possible, the quasi-degenerate scalar Higgs singlet produces a screening effect. Suppression of the branching ratio below the SM prediction is limited to at most a factor of $\sim 1/2$ at leading order.

The situation in the NMSSM is different. The CP-even singlet remains heavy close to the R-symmetric limit. The contribution of the singlet pseudoscalar can therefore easily exceed the experimental limits at large values of $\tan\beta$, unless the double-singlet mixing is constrained to be small. Due to the absence of an accompanying CP-even Higgs boson, the cancellation between SUSY and SM-like contribution to the branching ratio $B(B_s \to \mu^+\mu^-)$ can be highly efficient in this model, where suppression can be more than one order of magnitude relative to the SM prediction. Such an effect is not possible within the MSSM at large $\tan\beta$. Possible discovery of SUSY partners at the LHC along with non-observation of the decay $B_s \to \mu^+\mu^-$ would strongly point towards the NMSSM and searches for light pseudoscalar particles should become a high priority, should such a situation present itself in the near future.

The present work was carried out within the framework of minimal flavour violation. However, within a general renormalisable framework of the models we have been studying here, new sources of CP violation could be present amongst the soft SUSY-breaking terms. This is expected to lead to new CP-violating threshold corrections for both the singlet and doublet Higgs fields. Additionally, both the NMSSM and the MNSSM exhibit explicit CP violation in the tree-level Higgs potential. These effects may lead to significant enhancements of the signals considered here, along with contributions to other observables, such as electric dipole moments. We plan to report progress on these issues in the near future.

Acknowledgments

This work was partially supported by the STFC research grant: PP/D000157/1.

A The Chargino-Neutralino Propagator Matrix

In (2.2), we need to evaluate the different left- and right-handed chiral components of the chargino-neutralino propagator matrix:

$$\Delta_C(\not{k}) \;=\; \frac{1}{\not{k}\mathbf{1}_9 - \mathbf{M}_C P_L - \mathbf{M}_C^\dagger P_R} \;. \tag{A.1}$$

To do so, it proves more convenient to use the chiral representation for (A.1) and write $\Delta_C(\not{k})$ as

$$\Delta_C(\not{k}) \;=\; \left(\begin{array}{cc} -\mathbf{M}_C\mathbf{1}_2 & \mathbf{1}_9\mathbf{k} \\ \mathbf{1}_9\bar{\mathbf{k}} & -\mathbf{M}_C^\dagger\mathbf{1}_2 \end{array} \right)^{-1} \;. \tag{A.2}$$

In the above, we have defined $\mathbf{k} = k_\mu\sigma^\mu$ and $\bar{\mathbf{k}} = k_\mu\bar{\sigma}^\mu$, where $\sigma^\mu = (\mathbf{1}_2,\, \boldsymbol{\sigma})$, $\bar{\sigma}^\mu = (\mathbf{1}_2,\, -\boldsymbol{\sigma})$ and $\boldsymbol{\sigma}$ are the usual Pauli matrices. Note that $\mathbf{k}\bar{\mathbf{k}} = \bar{\mathbf{k}}\mathbf{k} = k^2\mathbf{1_2}$.

Our calculational task is equivalent to finding the inverse of an invertible, block matrix M,

$$M = \left(\begin{array}{cc} A & B \\ C & D \end{array} \right) \;, \tag{A.3}$$

where A, B, C and D are $m \times m$-, $m \times n$-, $n \times m$- and $n \times n$-dimensional matrices, respectively. To this end, we denote the inverse of M by

$$M^{-1} = \left(\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array} \right) \;. \tag{A.4}$$

Correspondingly, $\alpha,\ \beta,\ \gamma$ and δ are matrices of dimensionality $m \times m$, $m \times n$, $n \times m$ and $n \times n$. We may now use the condition $MM^{-1} = \mathbf{1}_{(n+m)}$ to write the four constraints,

$$A\alpha + B\gamma \;=\; \mathbf{1}_m \;, \tag{A.5}$$

$$C\alpha + D\gamma \;=\; \mathbf{0}_{n\times m} \;, \tag{A.6}$$

$$A\beta + B\delta \;=\; \mathbf{0}_{m\times n} \;, \tag{A.7}$$

$$C\beta + D\delta \;=\; \mathbf{1}_n \;. \tag{A.8}$$

Conditions (A.6) and (A.7) imply that

$$\gamma \;=\; -D^{-1}C\alpha \;, \tag{A.9}$$

$$\beta \;=\; -A^{-1}B\delta \;. \tag{A.10}$$

Combining these with (A.5) and (A.8), we may derive

$$\alpha \;=\; \Big(A - BD^{-1}C\Big)^{-1} \;, \tag{A.11}$$

$$\delta \;=\; \Big(D - CA^{-1}B\Big)^{-1} \;, \tag{A.12}$$

so that the inverse matrix M^{-1} may be written as

$$M^{-1} \;=\; \begin{pmatrix} (A - BD^{-1}C)^{-1} & -A^{-1}B\,(D - CA^{-1}B)^{-1} \\ -D^{-1}C\,(A - BD^{-1}C)^{-1} & (D - CA^{-1}B)^{-1} \end{pmatrix} . \tag{A.13}$$

One should observe here that the expression M^{-1} is not unique. Specifically, we could have used $M^{-1}M = \mathbf{1}_{m+n}$ to obtain equations analogous to (A.5)–(A.8) and express M^{-1} in a form which, although less compact, relies on the inverses of two combinations of submatrices only, e.g. D^{-1} and $(A - BD^{-1}C)^{-1}$. Such expressions are more efficiently implemented in numerical routines. However, all these alternative forms are equivalent to each other.

We now return to $\Delta_C(\not{k})$ in (A.2). Using (A.13), the chargino-neutralino propagator matrix may now be written down as

$$\Delta_C(\not{k}) \;=\; \begin{pmatrix} \mathbf{M}_C^\dagger \left(k^2\mathbf{1}_9 - \mathbf{M}_C\mathbf{M}_C^\dagger\right)^{-1}\mathbf{1}_2 & \left(k^2\mathbf{1}_9 - \mathbf{M}_C^\dagger\mathbf{M}_C\right)^{-1}\mathbf{k} \\ \left(k^2\mathbf{1}_9 - \mathbf{M}_C\mathbf{M}_C^\dagger\right)^{-1}\bar{\mathbf{k}} & \mathbf{M}_C\left(k^2\mathbf{1}_9 - \mathbf{M}_C^\dagger\mathbf{M}_C\right)^{-1}\mathbf{1}_2 \end{pmatrix} , \tag{A.14}$$

where we have used the fact that the only non-zero commutator amongst the submatrices is $[A, D]$ which simplifies the above expression. From (A.14), we may now obtain the different block elements, by appropriately acting with left- and right-handed projection operators $P_{L.R}$. For example, for the expression that occurs in (2.2), we find that

$$P_L\,\Delta_C(\not{k})\,P_L \;=\; \mathbf{M}_C^\dagger\left(k^2\mathbf{1}_9 - \mathbf{M}_C\mathbf{M}_C^\dagger\right)^{-1}\mathbf{1}_2 , \tag{A.15}$$

where we have suppressed the vanishing elements on the RHS of (A.15).

Using (A.15), we may numerically integrate expressions such as (2.2) without any need to first diagonalise the mass matrices appearing in the propagators. At each sample point in the integral, the propagator matrices are numerically inverted and the matrix element relevant to the process is selected, i.e. the propagator matrix element representing the transition between the required electroweak eigenstates. In this way, we avoid the introduction of cumbersome mixing matrices for the squarks and charginos in the analytic expressions of Section 2.

Furthermore, by working with all propagator matrices in the electroweak basis, the calculation of the derivatives appearing in (2.12) is greatly simplified. The expression for each derivate of the propagators in the matrix form contains no more than two terms. By contrast, if we were to diagonalise the mass matrices, we would find expressions containing three terms for each mass eigenstate (coming from applying the Leibniz rule to the product of Higgs-dependant masses with initial and final mixing matrices), many of which typically require further expansion before one could cast them into a useful compact form.

References

[1] S.L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. **D2** (1970) 1285.

[2] N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531;
M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.

[3] S. R. Choudhury and N. Gaur, Phys. Lett. B **451** (1999) 86 [arXiv:hep-ph/9810307];
K. S. Babu and C. F. Kolda, Phys. Rev. Lett. **84** (2000) 228 [arXiv:hep-ph/9909476];
C. S. Huang, W. Liao, Q. S. Yan and S. H. Zhu, Phys. Rev. D **63** (2001) 114021 [Erratum-ibid. D **64** (2001) 059902] [arXiv:hep-ph/0006250];
G. Isidori and A. Retico, JHEP **0111** (2001) 001 [arXiv:hep-ph/0110121];
A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **659** (2003) 3 [arXiv:hep-ph/0210145].

[4] C. Bobeth, T. Ewerth, F. Kruger and J. Urban, Phys. Rev. D **64** (2001) 074014 [arXiv:hep-ph/0104284];
A. Dedes, Mod. Phys. Lett. A **18** (2003) 2627 [arXiv:hep-ph/0309233].

[5] A. J. Buras, P. H. Chankowski, J. Rosiek and L. Slawianowska, Nucl. Phys. B **619** (2001) 434 [arXiv:hep-ph/0107048].

[6] A. Dedes and A. Pilaftsis, Phys. Rev. D **67** (2003) 015012 [arXiv:hep-ph/0209306].

[7] J. Ellis, J. S. Lee and A. Pilaftsis, Phys. Rev. D **76** (2007) 115011 [arXiv:0708.2079 [hep-ph]].

[8] J. E. Kim and H. P. Nilles, Phys. Lett. B **138** (1984) 150;
L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. D **27**, 2359 (1983);
G. F. Giudice and A. Masiero, Phys. Lett. B **206**, 480 (1988);
E. J. Chun, J. E. Kim and H. P. Nilles, Nucl. Phys. B **370**, 105 (1992);
I. Antoniadis, E. Gava, K. S. Narain and T. R. Taylor, Nucl. Phys. B **432**, 187 (1994).

[9] E. Accomando *et al.*, arXiv:hep-ph/0608079.

[10] P. Fayet, Nucl. Phys. B **90** (1975) 104;
J. M. Frere, D. R. T. Jones and S. Raby, Nucl. Phys. B **222** (1983) 11;
J. P. Derendinger and C. A. Savoy, Nucl. Phys. B **237** (1984) 307;
J. R. Ellis, J. F. Gunion, H. E. Haber, L. Roszkowski and F. Zwirner, Phys. Rev. D **39** (1989) 844;
P. N. Pandita, Phys. Lett. B **318** (1993) 338;
S. F. King and P. L. White, Phys. Rev. D **52** (1995) 4183 [arXiv:hep-ph/9505326];

B. Ananthanarayan and P. N. Pandita, Phys. Lett. B **353** (1995) 70 [arXiv:hep-ph/9503323];
M. Bastero-Gil, C. Hugonie, S. F. King, D. P. Roy and S. Vempati, Phys. Lett. B **489**, 359 (2000) [arXiv:hep-ph/0006198];
D. J. Miller, R. Nevzorov and P. M. Zerwas, Nucl. Phys. B **681** (2004) 3 [arXiv:hep-ph/0304049];
U. Ellwanger, J. F. Gunion and C. Hugonie, JHEP **0507** (2005) 041 [arXiv:hep-ph/0503203].

[11] C. Panagiotakopoulos and A. Pilaftsis, Phys. Rev. D **63** (2001) 055003 [arXiv:hep-ph/0008268];
A. Dedes, C. Hugonie, S. Moretti and K. Tamvakis, Phys. Rev. D **63** (2001) 055009 [arXiv:hep-ph/0009125];
A. Menon, D. E. Morrissey and C. E. M. Wagner, Phys. Rev. D **70** (2004) 035005 [arXiv:hep-ph/0404184];
S. W. Ham, S. K. Oh, C. M. Kim, E. J. Yoo and D. Son, Phys. Rev. D **70** (2004) 075001 [arXiv:hep-ph/0406062];
C. Balazs, M. S. Carena, A. Freitas and C. E. M. Wagner, JHEP **0706** (2007) 066 [arXiv:0705.0431 [hep-ph]];
S. Hesselbach, D. J. Miller, G. Moortgat-Pick, R. Nevzorov and M. Trusov, Phys. Lett. B **662** (2008) 199 [arXiv:0712.2001 [hep-ph]].

[12] U. Ellwanger, J. F. Gunion, C. Hugonie and S. Moretti, arXiv:hep-ph/0305109;
S. Chang, P. J. Fox and N. Weiner, JHEP **0608** (2006) 068 [arXiv:hep-ph/0511250];
K. Cheung, J. Song and Q. S. Yan, Phys. Rev. Lett. **99** (2007) 031801 [arXiv:hep-ph/0703149];
M. Carena, T. Han, G. Y. Huang and C. E. M. Wagner, JHEP **0804** (2008) 092 [arXiv:0712.2466 [hep-ph]];
J. R. Forshaw, J. F. Gunion, L. Hodgkinson, A. Papaefstathiou and A. D. Pilkington, JHEP **0804** (2008) 090 [arXiv:0712.3510 [hep-ph]].

[13] A. Arhrib, K. Cheung, T. J. Hou and K. W. Song, JHEP **0703** (2007) 073 [arXiv:hep-ph/0606114].

[14] M. A. Sanchis-Lozano, J. Phys. Soc. Jap. **76** (2007) 044101 [arXiv:hep-ph/0610046];
R. Dermisek, J. F. Gunion and B. McElrath, Phys. Rev. D **76** (2007) 051105(R) [arXiv:hep-ph/0612031];
E. Fullana and M. A. Sanchis-Lozano, Phys. Lett. B **653** (2007) 67 [arXiv:hep-ph/0702190];
R. N. Hodgkinson, Phys. Lett. B **665** (2008) 219 [arXiv:0802.3197 [hep-ph]].

[15] T. Banks, Nucl. Phys. B **303** (1988) 172;
E. Ma, Phys. Rev. D **39** (1989) 1922.

[16] R. Hempfling, Phys. Rev. D **49** (1994) 6168;
L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50** (1994) 7048 [arXiv:hep-ph/9306309];
M. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, Nucl. Phys. B **426** (1994) 269 [arXiv:hep-ph/9402253];
D. M. Pierce, J. A. Bagger, K. T. Matchev and R. j. Zhang, Nucl. Phys. B **491** (1997) 3 [arXiv:hep-ph/9606211];
F. Borzumati, G. R. Farrar, N. Polonsky and S. D. Thomas, Nucl. Phys. B **555** (1999) 53 [arXiv:hep-ph/9902443].

[17] J. A. Coarasa, R. A. Jimenez and J. Sola, Phys. Lett. B **389** (1996) 312 [arXiv:hep-ph/9511402];
R. A. Jimenez and J. Sola, Phys. Lett. B **389** (1996) 53 [arXiv:hep-ph/9511292];
K. T. Matchev and D. M. Pierce, Phys. Lett. B **445** (1999) 331 [arXiv:hep-ph/9805275];
P. H. Chankowski, J. R. Ellis, M. Olechowski and S. Pokorski, Nucl. Phys. B **544** (1999) 39 [arXiv:hep-ph/9808275];
K. S. Babu and C. F. Kolda, Phys. Lett. B **451** (1999) 77 [arXiv:hep-ph/9811308].

[18] R. N. Hodgkinson and A. Pilaftsis, Phys. Rev. D **76** (2007) 015007 [arXiv:hep-ph/0612188].

[19] G. Hiller, Phys. Rev. D **70** (2004) 034018 [arXiv:hep-ph/0404220].

[20] Z. Heng, R. J. Oakes, W. Wang, Z. Xiong and J. M. Yang, Phys. Rev. D **77** (2008) 095012 [arXiv:0801.1169 [hep-ph]].

[21] S. R. Coleman and E. Weinberg, Phys. Rev. D **7** (1973) 1888.

[22] A. Lenz and U. Nierste, JHEP **0706** (2007) 072 [arXiv:hep-ph/0612167].

[23] A. J. Buras, S. Jager and J. Urban, Nucl. Phys. B **605** (2001) 600 [arXiv:hep-ph/0102316];
M. Ciuchini *et al.*, JHEP **0107** (2001) 013 [arXiv:hep-ph/0012308];
A. J. Buras, M. Misiak and J. Urban, Nucl. Phys. B **586** (2000) 397 [arXiv:hep-ph/0005183];
D. Becirevic, V. Gimenez, G. Martinelli, M. Papinutto and J. Reyes, Nucl. Phys. Proc. Suppl. **106** (2002) 385 [arXiv:hep-lat/0110117].

[24] R. Arnowitt, B. Dutta, T. Kamon and M. Tanaka, Phys. Lett. B **538** (2002) 121 [arXiv:hep-ph/0203069];
S. Baek, P. Ko and W. Y. Song, JHEP **0303** (2003) 054 [arXiv:hep-ph/0208112].

[25] J. S. Lee, A. Pilaftsis, M. S. Carena, S. Y. Choi, M. Drees, J. R. Ellis and C. E. M. Wagner, Comput. Phys. Commun. **156** (2004) 283 [arXiv:hep-ph/0307377];
J. S. Lee, M. Carena, J. Ellis, A. Pilaftsis and C. E. M. Wagner, arXiv:0712.2360 [hep-ph].

[26] G. Buchalla and A. J. Buras, Nucl. Phys. B **412** (1994) 106 [arXiv:hep-ph/9308272];
M. Misiak and J. Urban, Phys. Lett. B **451** (1999) 161 [arXiv:hep-ph/9901278].

[27] B. A. Dobrescu, G. L. Landsberg and K. T. Matchev, Phys. Rev. D **63** (2001) 075003 [arXiv:hep-ph/0005308];
B. A. Dobrescu and K. T. Matchev, JHEP **0009** (2000) 031 [arXiv:hep-ph/0008192].